

ІНСТИТУТ
ФІЗИКИ
КОНДЕНСОВАНИХ
СИСТЕМ

ICMP-00-18E

Maxym Dudka, Reinhard Folk*, Yurij Holovatch

PHASE TRANSITION IN THE RANDOM ANISOTROPY
MODEL

*Institut für theoretische Physik, Johannes Kepler Universität Linz, A-4040
Linz, Austria

УДК: 530.145

PACS: 61.43.-j,64.60.Ak

Фазовий перехід в моделі з випадковою анізотропією

М. Дудка, Р. Фольк, Ю. Головач

Анотація. Досліджується вплив локальної анізотропії випадкової орієнтації на ферромагнітний фазовий перехід для двох випадків розподілу осей анізотропії. З цією метою за допомогою методу теоретико-польової ренормалізаційної групи проведено аналіз моделі магнетика з випадковою анізотропією в двопетевому наближенні уточненому пересумовуванням асимптотичних рядів. Підтверджено однопетлевий результат Агароні, який вказує на відсутність фазового переходу другого роду для ізотропного розподілу осей випадкової анізотропії при вимірності простору $d < 4$. Для кубічного розподілу досяжна стійка точка приводить до критичних показників розведеного Ізинга.

Phase transition in the random anisotropy model

Maxym Dudka, Reinhard Folk, Yurij Holovatch

Abstract. The influence of a local anisotropy of random orientation on a ferromagnetic phase transition is studied for two cases of anisotropy axis distribution. To this end a model of a random anisotropy magnet is analysed by means of a field theoretical renormalization group approach in two loop approximation refined by resummation of asymptotic series. The one-loop result of Aharony indicating the absence of a 2nd order phase transition for isotropic distribution of random anisotropy axis at space dimension $d < 4$ is corroborated. For a cubic distribution the accessible stable fixed point leads to disordered Ising-like critical exponents.

Подається в Fluctuating Partns and Fields (World Scientific, Singapore, 2001)

Submitted to Fluctuating Partns and Fields (World Scientific, Singapore, 2001)

1. Introduction

Modern understanding of universal properties of matter in the vicinity of critical points to a great extent is due to the application of the renormalization group (RG) ideas [1]. Applied in the problems of condensed matter physics in the early 70-ies the RG technique proved to be a powerful tool to study critical phenomena. For example, expressions for critical exponents governing magnetic phase transition in regular systems are known by now with record accuracy both for isotropic [2] ($O(m)$ symmetrical) and cubic [3] magnets. The RG approach also sheds light on the influence of structural disorder on ferromagnetism. In the present paper we will apply the field theoretical RG approach to study peculiarities of magnetic behaviour influenced by disorder in a form of random anisotropy axis [4]. It is special pleasure for us to dedicate this paper to Prof. Hagen Kleinert on the occasion of his 60th anniversary. His contribution to the field is hard to be overestimated.

Although an influence of a weak quenched structural disorder on universal properties of a ferromagnetic phase transition is a problem of intensive study during already several decades there remains a number of unsettled questions. Here, one should discriminate between random site, random-field and random anisotropy magnets. A weak quenched disorder preserves 2nd order phase transition in three dimensional ($d = 3$) random site magnets [5] but can destroy this transition in random field systems [6] for $d < 4$. The situation for the random-anisotropy magnets is not so clear.

Typical examples of random-anisotropy magnets are amorphous rare-earth – transition metal alloys. Some of these systems order magnetically and for the description of the ordered structure it has been proposed [4] to consider a regular lattice of magnetic ions, each of them being a subject to a local anisotropy of random orientation. The Hamiltonian of this random anisotropy model (RAM) reads [4]:

$$\mathcal{H} = - \sum_{\mathbf{R}, \mathbf{R}'} J_{\mathbf{R}, \mathbf{R}'} \vec{S}_{\mathbf{R}} \vec{S}_{\mathbf{R}'} - D_0 \sum_{\mathbf{R}} (\hat{x}_{\mathbf{R}} \vec{S}_{\mathbf{R}})^2, \quad (1)$$

where $\vec{S}_{\mathbf{R}}$ is an m -component vector on a lattice site \mathbf{R} , $J_{\mathbf{R}, \mathbf{R}'}$ is an exchange interaction, D_0 is an anisotropy strength, and $\hat{x}_{\mathbf{R}}$ is a unit vector pointing in the local (quenched) random direction of an uniaxial anisotropy.

The model was investigated by variety of techniques including mean-field theory [7], computer simulations [8], $1/m$ -expansion [9], renormalization group ε -expansion [10–12]. The limiting case of an infinite

anisotropy was subject of a detailed study as well [13,14]. However the nature of low-temperature phase in RAM is not completely clear up to now, and several low-temperature phases were discussed like ferromagnetic ordering [7,8], spin-glass phase [8,9], and quasi long-range ordering [15].

The nature of ordering is connected with the distribution of the random variables $\hat{x}_{\mathbf{R}}$ in (1). For an isotropic distribution arguments similar to those applied by Imry and Ma [16] for a random-field Ising model bring about the absence of ferromagnetic order for space dimensions $d < 4$ [12,17]. Whereas anisotropic distributions may lead to a ferromagnetic order [18].

Application of Wilson RG technique to RAM with the isotropic distribution of a local anisotropy axis suggests [10] the possibility of “runaway” solutions of the recursion equations. Such behaviour was interpreted as a smeared transition. However this result was obtained in first order of ε -expansion and remains to be confirmed also in higher orders.

Here, we will report results obtained by means of the field theoretical RG technique in two loop approximation refined by resummation of the resulting asymptotic series. We will consider two cases of distribution of the random anisotropy axis and show that a ferromagnetic second order phase transition takes place only when the distribution is non-isotropic. Moreover we will show that the RAM provides one more example of a disordered model, where the only possible new critical behaviour is of “random Ising” type, similar to the site-diluted magnets [5]. More detailed results can be found in Refs [19,20].

2. Isotropic case

In order to deal with quenched disorder one way to obtain the effective Hamiltonian of a RAM is to make use of the replica trick. Then for a given configuration of quenched random variables $\hat{x}_{\mathbf{R}}$ in (1) the partition function may be written in a form of functional integral of a Gibbs distribution depending on $\hat{x}_{\mathbf{R}}$. To perform averaging over configurations one should complete the model by choosing certain distribution of $\hat{x}_{\mathbf{R}}$. We will analyse two cases: when the random vector \hat{x} points with equal probability in any direction of the m -dimensional hyperspace (isotropic case) and when \hat{x} lies along the edges of m -dimensional hypercube (cubic case). Other distributions may be considered as well. For the first case the distribution function reads:

$$p(\hat{x}) \equiv \left(\int d^m \hat{x} \right)^{-1} = \frac{\Gamma(m/2)}{2\pi^{m/2}}. \quad (2)$$

Following the above described program one ends up with the replica $n \rightarrow 0$ limit of the effective Hamiltonian [10]:

$$\mathcal{H}_{eff} = - \int d^d R \left\{ \frac{1}{2} \left[\mu_0^2 |\vec{\varphi}|^2 + |\vec{\nabla} \vec{\varphi}|^2 \right] + u_0 |\vec{\varphi}|^4 + v_0 \sum_{\alpha=1}^n |\vec{\varphi}^\alpha|^4 + w_0 \sum_{\alpha,\beta=1}^n \sum_{i,j=1}^m \phi_i^\alpha \phi_j^\alpha \phi_i^\beta \phi_j^\beta \right\}, \quad (3)$$

here, μ_0^2 and u_0, v_0, w_0 are defined by D_0 and familiar bare couplings of an m -vector model, and $\vec{\varphi}^\alpha \equiv \vec{\phi}_{\mathbf{R}}^\alpha$ is a m -dimensional vector, $|\vec{\varphi}|^2 = \sum_{\alpha=1}^n |\vec{\varphi}^\alpha|^2$. Bare couplings $u_0 > 0, v_0 > 0, w_0 < 0$. Furthermore, values of u_0 and w_0 are related to appropriate cumulants of the distribution function (2) and their ratio equals $w_0/u_0 = -m$. Note that the symmetry of u_0 and v_0 terms corresponds to the random site m -vector model [21]. However the u_0 -term has an opposite sign.

In order to study long-distance properties of the Hamiltonian (3), we use the field-theoretical RG approach [1]. In this approach the critical point of a system corresponds to a stable fixed point (FP) of the RG transformation. We apply the massive field theory renormalization scheme [22] performing renormalization at fixed space dimension d and zero external momenta. In two loop approximation we get [19] expressions for the RG functions in form of asymptotic series in renormalized couplings u, v, w .

As it was mentioned in the introduction the only known RG results for RAM with isotropic distribution of the local anisotropy axis so far are those obtained in first order in ε [10]. In particular one gets eight fixed points. All FPs with $u > 0, v > 0, w < 0$ appear to be unstable for $\varepsilon > 0$ except of the ‘‘polymer’’ $O(n=0)$ FP III which is stable for all m (see Fig. 1). However the presence of a stable FP is not a sufficient condition for a 2nd order phase transition. In order to be physically relevant the FP should be accessible from the initial values of couplings. This is not the case for the location of FPs shown in Fig. 1. Indeed starting from the region of physical initial conditions (denoted by the cross in the Fig. 1) in the plane of $v = 0$ one would have to cross the separatrix joining the unstable FPs I and VI. This is not possible and so one never reaches the stable FP III. As far as both FPs I and VI are strongly unstable with respect to v FP III is not accessible for arbitrary positive v either. Finally one ends up with the conclusion about absence of the 2nd order phase transition in the model as runaway solutions of the RG equations show.




Figure 1. Fixed points of the RAM with isotropic distribution of a local anisotropy axis. The only fixed points located in the octant $u > 0, v > 0, w < 0$ are shown. The filled box shows the stable fixed point, the cross denotes typical initial values of couplings.

FP	m	u^*	v^*	w^*	ν	η
I	$\forall m$	0	0	0		
II	2	0	0.9107	0	0.663	0.027
	3	0	0.8102	0	0.693	0.027
	4	0	0.7275	0	0.724	0.027
III	$\forall m$	1.1857	0	0	0.590	0.023
IV	2	-0.0322	0.9454	0	0.668	0.027
	3	0.1733	0.6460	0	0.659	0.027
	4	0.2867	0.4851	0	0.653	0.028
VI	2	1.4650	0	-1.6278	0.449	-0.028
VIII	2	0.7517	0.7072	-0.3984	0.626	0.031
	3	0.8031	0.5463	-0.3305	0.620	0.029
	4	0.8349	0.4545	-0.2888	0.617	0.029

Table 1. Resummed values of the fixed points and critical exponents for isotropic case in two-loop approximation for $d = 3$.

The main question of interest here is whether the above described picture of runaway solutions is not an artifact of an ε -expansion. To check this we used a more refined analysis of the FPs and their stability, considering the series for RG functions directly at $d = 3$ [22]. As it is known series of this type are asymptotic at best and a resummation procedure has to be applied to obtain reliable data on their basis. We made use of Padé–Borel resummation techniques [23] first writing the RG functions as resolvent series [24] in one auxiliary variable and then performing resummation. Numerical values of the FPs are given in the Table 1 [25]. Resummed two-loop results qualitatively confirm the picture obtained in the first order in ε -expansion: stability of the FPs does not change after resummation. This supports the conjecture of Aharony [10] about absence of accessible stable FP for the RAM with isotropic distribution of the local anisotropy axis. We display in the table resummed in a similar way values of the correlation length and pair correlation function critical exponents ν and η : being calculated in unstable FPs they are rather to be considered as effective ones.

3. Cubic case

Let us now consider the second example of anisotropy axis distribution, when the vector $\hat{x}_{\mathbf{R}}$ (1) points only along one of the $2m$ directions of axes \hat{k}_i of a cubic lattice :

$$p(\hat{x}) = \frac{1}{2m} \sum_{i=1}^m [\delta^{(m)}(\hat{x} - \hat{k}_i) + \delta^{(m)}(\hat{x} + \hat{k}_i)]. \quad (4)$$

The *rationale* for such a choice is to mimic the situation when an amorphous magnet still “remembers” initial (cubic) lattice structure. Repeating the procedure described in the previous section one ends up with the following effective Hamiltonian (being of interest in the limit $n \rightarrow 0$) [10]:

$$\mathcal{H}_{eff} = - \int d^d R \left\{ \frac{1}{2} \left[\mu_0^2 |\vec{\varphi}|^2 + |\vec{\nabla} \vec{\varphi}|^2 \right] + u_0 |\vec{\varphi}|^4 + v_0 \sum_{\alpha=1}^n |\vec{\phi}^\alpha|^4 + w_0 \sum_{i=1}^m \sum_{\alpha,\beta=1}^n \phi_i^{\alpha 2} \phi_i^{\beta 2} + y_0 \sum_{i=1}^m \sum_{\alpha=1}^n \phi_i^{\alpha 4} \right\}. \quad (5)$$

Here, the bare couplings are $u_0 > 0$, $v_0 > 0$, $w_0 < 0$. The y_0 term is generated when the RG transformation is applied and may be of either

sign. The symmetry of w_0 terms differs in (3) and (5). Furthermore, values of u_0 and w_0 differ for Hamiltonians (3) and (5) but their ratio equals $-m$ again.

We apply the massive field theory renormalization scheme [22] and get the RG functions in two loop approximation [20]. As well as in the previous case we reproduce the first-order ε -results [10]. Now one gets 14 FPs. However, in first order of ε -expansion all FPs with $u > 0$, $v > 0$, $w < 0$ appear to be unstable for $\varepsilon > 0$ except of the “polymer” $O(n = 0)$ FP III which is stable for all m but not accessible (see Fig. 2). Now the account of the ε^2 terms qualitatively changes the picture: indeed, the system of equations for the FPs appears to be degenerated at the one loop level. As known from other cases in two loop order this leads to the appearance of a new FP which is stable and is expressed by a $\sqrt{\varepsilon}$ series [21]. The possibility of such scenario was predicted already in Refs [18]. However it remained unclear whether there exist any other accessible stable FPs.




Figure 2. Fixed points of the RAM with distribution of a local anisotropy axis along hypercube axis for $v = 0$. The only fixed points located in the region $u > 0$, $w < 0$ are shown. Filled boxes show the stable fixed points, the cross denotes typical initial values of couplings.

Applying Padé–Borel resummation we get 16 FPs. Values of the FPs with $u^* > 0$, $v^* > 0$, $w^* < 0$ are presented in the Table 2 [25]. The last FP XV in Table 2 corresponds the stable FP of $\sqrt{\varepsilon}$ -expansion. It has coordinates with $u^* = v^* = 0$, $w^* < 0$ and $y^* > 0$ and is accessible

from the typical initial values of couplings (shown by cross in the Fig.2). Applying the resummation procedure we did not find any other stable FPs in the region of interest. The effective Hamiltonian (5) at $u = v = 0$ in the replica limit $n \rightarrow 0$ reduces to a product of m effective Hamiltonians of a weakly diluted quenched random site Ising model. This means that for any value of $m > 1$ the system is characterized by the same set of critical exponents as those of a weakly diluted random site quenched Ising model. We give in the Table 2 values of critical exponents in the other FPs as well: if the flows from initial values of couplings pass near the these FPs one may observe an effective critical behaviour governed by these critical exponents.

FP	m	u^*	v^*	w^*	y^*	ν	η
I	$\forall m$	0	0	0	0	1/2	0
II	2	0	0.9107	0	0	0.663	0.027
	3	0	0.8102	0	0	0.693	0.027
	4	0	0.7275	0	0	0.720	0.026
III	$\forall m$	1.1857	0	0	0	0.590	0.023
V	$\forall m$	0	0	0	1.0339	0.628	0.026
VI	3	0.1733	0.6460	0	0	0.659	0.027
	4	0.2867	0.4851	0	0	0.653	0.027
VII	$\forall m$	2.1112	0	-2.1112	0	1/2	0
VIII	2	0	1.5508	0	-1.0339	0.628	0.026
	3	0	0.8393	0	-0.0485	0.693	0.027
	4	0	0.5259	0	0.3624	0.709	0.026
IX	3	0.1695	0.7096	0	-0.1022	0.659	0.027
	4	0.2751	0.4190	0	0.1432	0.653	0.027
X	$\forall m$	0.6678	0	-0.6678	1.0339	0.628	0.026
XV	$\forall m$	0	0	-0.4401	1.5933	0.676	0.031

Table 2. Resummed values of the FPs and critical exponents for cubic distribution in two-loop approximation for $d = 3$.

4. Conclusions

In this paper we applied the field theoretical RG approach to analyze critical behaviour of a model of random anisotropy magnet with isotropic and cubic distributions of a local anisotropy axis. The origin of a low-temperature phase in this model is not completely clear. General ar-

guments based on an estimate of the energy for formation of magnetic domains [16] lead to the conclusion that for $d < 4$ a ferromagnetic order is absent [12,17]. However these arguments do not take into account entropy which may be important for disordered systems [14]. Furthermore, these arguments do not apply for anisotropic distributions of the random axis [18].

In the RG analysis absence of a ferromagnetic second order phase transition corresponds to the lack of a stable FP of the RG transformation. However in the case of RAM with isotropic distribution of a local anisotropy axis the scenario differs. Our two loop calculation brings about the presence of a $O(n = 0)$ symmetric FP which is stable for any value of m for both isotropic and cubic distributions of a random anisotropy axis. However, this FP is not accessible from the initial values of the couplings. We checked the location of the FPs up to second order in ε -expansion and by means of a fixed $d = 3$ technique refined by Padé-Borel resummation.

In the case of isotropic distribution of a random anisotropy axis our analysis supports the conjecture of Aharony [10] based on linear in ε -results about runaway solutions of the RG equations. For the cubic distribution we get two stable FPs. One of them (FP III in Fig. 2) is not accessible as in the isotropic case. But the disordered Ising-like FP (FP XV in Fig. 2) may be reached from the initial values of couplings. Applying the resummation procedure we did not find any other stable FPs in the region of interest. This means that RAM with cubic distributions of a random anisotropy axis is governed by a set of critical exponents of a weakly site diluted quenched Ising model [21,26].

To conclude we want to attract attention to certain similarity in the critical behaviour of both random-site [21] and random-anisotropy [4] quenched magnets: if at all there appears *new critical behaviour* it *always is governed by critical exponents of site-diluted Ising type*. The above calculations of a “phase diagram” of RAM were based on two loop expansions improved by a resummation technique. Once the qualitative picture became clear there is no need to go into higher orders of a perturbation theory as far as the critical exponents of site-diluted Ising model are known by now with high accuracy [26].

Yu. H. acknowledges helpful discussions with Mykola Shpot. This work has been supported in part by “Österreichische Nationalbank Jubiläumsfonds” through the grant No 7694.

References

1. D.J. Amit *Field Theory, the Renormalization Group, and Critical Phenomena*. (World Scientific, Singapore, 1984); J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena* (Oxford University Press, Oxford, 1989); H. Kleinert and V. Schulte-Frohlinde, *Critical Properties of ϕ^4 -Theories*.(World Scientific , Singapore, 2000).
2. H. Kleinert *et al*, Phys. Lett. B, **272**, 39, (1991), Phys. Lett. B, **319**, 545, (1993); R. Guida and J. Zinn-Justin, J. Phys. A, **31**, 8103,(1998).
3. H. Kleinert and V. Schulte-Frohlinde, Phys. Lett. B, **342**, 284, (1995); J.M. Carmona, A. Pelissetto, and E. Vicari, Phys. Rev. B, **61**, 15136, (2000).
4. R. Harris, M. Plischke, and M.J. Zuckermann, Phys. Rev. Lett., **31**, 160, (1973).
5. A weak quenched random site dilution is irrelevant if the heat capacity critical exponent α_{pure} of the undiluted system is negative (A.B. Harris, J. Phys. C, **7**, 1671, (1974)). For positive α_{pure} dilution causes change in the universality class (see e.g. R. Folk, Yu. Holovatch, and T. Yavors’kii, Phys. Rev. B, **61**, 15114, (2000) for a recent review).
6. See e.g. reviews: D.P. Belanger and A.P. Young (JMMM, **100**, 272, (1991)); D.P. Belanger, eprint:cond-mat/0009029.
7. B. Derrida and J. Vannimenus, J. Phys. C, **13**, 3261, (1980); R. Harris and D. Zobin, J. Phys. F, **7**, 337, (1977); E. Callen, Y.J. Liu, and J.R. Cullen, Phys. Rev. B, **16**, 263, (1977); J.D. Patterson, G.R. Gruzalski, and D.J. Sellmyer, Phys. Rev. B, **18**, 1377, (1978); R. Alben, J.J. Becker, and M.C. Chi, J. Appl.Phys., **49**, 1653, (1978).
8. M.C. Chi and R. Alben, J. Appl. Phys., **48**, 2987, (1977); M.C. Chi and T. Egami, J. Appl. Phys., **50**, 165, (1979); R. Harris and S.H. Sung, J. Phys. F, **8**, L299, (1978); C. Jayaprakash and S. Kirkpatrick, Phys. Rev. B, **21**, 4072, (1980).
9. Y.Y. Goldschmidt, Nucl. Phys. B, **225**, 123, (1982); Y.Y. Goldschmidt, Phys. Rev. B, **30**, 1632, (1984); A. Khurana, A. Jagannathan, and M.J. Kosterlitz, Nucl. Phys. B, **40**, 1, (1984); A. Jagannathan, B. Shaub, and M.J. Kosterlitz, Nucl. Phys. B, **265**, 324, (1986); Y.Y. Goldschmidt and A. Aharony, Phys. Rev. B, **32**, 264, (1985).
10. A. Aharony, Phys. Rev. B, **12**, 1038, (1975).
11. J.H. Chen and T.C. Lubensky, Phys. Rev. B, **16**, 1206, (1977).
12. R.A. Pelcovits, E. Pytte, and J. Rudnick, Phys. Rev. Lett., **40**, 476, (1978); R.A. Pelcovits, *Phys. Rev. B***19**, 465, (1979); E. Pytte, Phys. Rev. B, **18**, 5046, (1978).
13. A. Bray and M. Moore, J. Phys. C, **18**, L139, (1985); A.B. Harris, R.G. Caflisch and J.R. Banavar, Phys. Rev. B, **35**, 4929, (1987).
14. K.M. Fischer and A. Zippelius, *J. Phys. C*, **18**, L1139, (1985).
15. A. Aharony and E. Pytte, Phys. Rev. Lett., **19**, 1583, (1980); R. Fisch, Phys. Rev. B, **57**, 269, (1998); D.E. Feldman, JETP Lett., **70**, 135, (1999); J. Chakrabaty, Phys. Rev. Lett. **81**, 385, (1998).
16. Y. Imry and S.-k. Ma, Phys. Rev. Lett., **35**, 1399, (1975).
17. S.-k. Ma and J. Rudnick, Phys. Rev. Lett., **40**, 589, (1978).
18. D. Mukamel and G. Grinstein, Phys. Rev. B, **25**, 381, (1982); A.L. Korzhenevskii and A.A. Luzhkov, Sov. Phys. JETP, **94**, 250, (1988).
19. M. Dudka, R. Folk, and Yu. Holovatch, Cond. Matt. Phys., **4**, (2001) (to appear)
20. M. Dudka, R. Folk, and Yu. Holovatch, in preparation
21. G. Grinstein and A. Luther, Phys. Rev. B, **13**, 1329, (1976).
22. G. Parisi (1973) (unpublished); J. Stat. Phys., **23**, 49, (1980).
23. G.A. Jr. Baker, B.G. Nickel, and D.I. Meiron, Phys. Rev. B **17**, 1365, (1978).
24. P.J.S. Watson, J. Phys. A., **7**, L167, (1974).
25. We absorb the value of a one-loop integral into the normalization of couplings.
26. Up to now the critical exponents of a weakly diluted quenched Ising model are calculated in RG technique with record six-loop accuracy in massive theory scheme: A. Pelissetto, and E. Vicari, eprint: cond-mat/0002402. In minimal subtraction scheme five loop results are available: B.N. Shalaev, S.A. Antonenko, and A.I. Sokolov, Phys. Lett. A, **230**, 105, (1997); R. Folk, Yu. Holovatch, and T. Yavorskii, J. Phys. Stud., **2**, 213, (1998); ZhETP Lett., **69**, 747, (1999).

Препринти Інституту фізики конденсованих систем НАН України розповсюджуються серед наукових та інформаційних установ. Вони також доступні по електронній комп'ютерній мережі на WWW-сервері інституту за адресою <http://www.icmp.lviv.ua/>

The preprints of the Institute for Condensed Matter Physics of the National Academy of Sciences of Ukraine are distributed to scientific and informational institutions. They also are available by computer network from Institute's WWW server (<http://www.icmp.lviv.ua/>)

Максим Леонідович Дудка
Райнгард Фольк
Юрій Васильович Головач

ФАЗОВИЙ ПЕРЕХІД В МОДЕЛІ З ВИПАДКОВОЮ АНІЗОТРОПІЄЮ

Роботу отримано 11 грудня 2000 р.

Затверджено до друку Вченою радою ІФКС НАН України

Рекомендовано до друку семінаром відділу СТеКС

Виготовлено при ІФКС НАН України

© Усі права застережені